

Standard Practice for Serving Protocol for Sensory Evaluation of Foods and Beverages¹

This standard is issued under the fixed designation E 1871; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This practice describes suggested procedures for presenting samples to sensory assessors. The purpose of this practice is to provide general guidelines for conducting sensory evaluation on a variety of foods and beverages, excluding beverage alcohol.

1.2 This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

2. Referenced Documents

2.1 ASTM Standards:

- E 1627 Practice for Sensory Evaluation of Edible Oils and Fats^2
- E 1810 Practice for Evaluating the Effects of Contaminants on the Odor and Waste of Exposed Fish²
- 2.2 IEC Standard:
- IEC 705 Guideline for Power Output Measurement of Consumer Microwave Ovens³

3. Summary of Practice

3.1 Consistency must be maintained in all aspects of preparation and serving of samples to ensure reproducible data. Guidelines for consistency may change with the test objective.

4. Significance and Use

4.1 This practice provides general guidelines and recommendations for presenting samples to assessors for sensory evaluation. Specific situations may require variations to these guidelines.

5. Procedure

5.1 General Guidelines:

5.1.1 *Pretest*—A practice session may be conducted with a few staff members or assessors to determine if the selected

procedures are appropriate for a specific test. Serving sizes, number of samples, number of questions, palate cleansers, sample temperature, lighting, etc., should be determined by pretesting and appropriately modified for the actual test.

5.1.2 *Product Variability*—Variability exists in all products. How product variability is handled depends on the objective of the test, the size of the effect one is attempting to detect, and the risks associated with decision making. Unless the test is designed to understand the extent of product variability, it should be minimized.

5.1.2.1 Samples must be consistent and uniform regarding production lot, age, package size, etc. All test conditions and serving procedures are determined by the test objective, test method, and test design. A descriptive test might have different requirements than an acceptance or preference test. It is important that all sample preparation and serving procedures be pretested.

5.1.2.2 Consider product variability when preparing samples for a test. Based on the objective of the test, contents of all containers may be blended for a given product prior to portioning individual samples or samples may be portioned from individual containers. For example, if six bottles of a given brand of salad dressing are required for a test, it may be decided to blend the contents of the six bottles. While this option appears to provide the most uniform sampling, it masks any variation among the six bottles. This can impact the results if a positive or negative effect goes undetected.

5.1.2.3 Another consideration is the pouring sequence for products that layer, for example, blended beverages and salad dressings. Each sample should be taken from the same location within the container for all variables for a single assessor. Another consideration of product variability is the difference between exposed surfaces and interior surfaces. Some products, for example, peanut butter, mayonnaise, butter, jam, and frozen desserts, may require removal of exposed surfaces prior to evaluation. It may be important, however, to evaluate both exposed and interior surfaces.

5.1.3 *Serving Containers/Utensils*—The same type of containers and utensils should be used throughout the test. Consider the following when choosing the container/utensil:

(*a*) The test objective, for example, a lid may be needed for aroma evaluation,

(b) Maintaining sample characteristics, for example, size, shape, serving temperature, moisture, etc.,

¹ This practice is under the jurisdiction of ASTM Committee E-18 on Sensory Evaluation of Materials and Products and is the direct responsibility of Subcommittee E18.06 on Food and Beverage Evaluation.

Current edition approved May 10, 1997. Published August 1997.

² Annual Book of ASTM Standards, Vol 15.07.

³ Available from International Microwave Power Institute, 13542 Union Village Circle, Clifton, VA 22024. Phone: (703) 830-5588.

(c) Product/container interaction,

 $\left(d\right)$ The amount of sample required to complete the evaluation,

(e) The amount of effort required by assessors to remove the product from the container for evaluation, and

(*f*) The containers/utensils should be neutral in color (unless tint is needed to mask color differences) and made of materials that are inert, nonreactive, and odor-free.

5.1.3.1 Some plastic materials are less inert, more susceptible to temperature changes, and less odor-free than others so they should be pretested prior to their use. Consider washing glassware with an unscented detergent, followed by baking at 93°C (200°F) for several hours, to eliminate any dusty smell. Assessors should not be able to draw conclusions about the identity of the samples due to extraneous variables not related to the test objective. It may be possible to mask color differences with light filters, subdued lighting, or tinted glassware, or combination thereof.

5.1.4 *Serving Size*—Consider the test objective when determining serving size. Provide enough of the sample to ensure the assessor evaluates the overall product, not just one or two components. The amount and size of the sample must be consistent for each assessor. Scoops, measuring cups, and top loading balances are effective for maintaining consistency.

5.1.4.1 Consider all of the test variables and product characteristics when determining sample preparation and serving protocol, for example, the top crust of bread is different from the bottom crust. Pretesting is helpful to accurately determine the amount needed, for example, the perceived spice level of a barbecue sauce may be markedly higher on the third or fourth bite than the first, so consider providing enough sauce for evaluation of both the first and third bites. It may be necessary to provide instructions to ensure that the samples are evaluated correctly, for example, "place entire sample in mouth," "be sure your bite includes the crust," etc.

5.1.4.2 It may be necessary for the assessors to eat an entire serving portion based on the test objective. Assessors should be informed of the quantity they will be required to consume prior to beginning the test. To ensure assessors comply, require them to finish the portion before completing the evaluation.

5.1.5 Serving Temperature-Consider the test objective in determining the sample serving temperature. The serving temperature of all samples must be consistent and within a predetermined range. The serving temperature, however, may be different than the evaluation temperature due to heat gain or loss. For example, a sample that is to be evaluated at 70°C may need to be served at 85°C. Since serving temperatures vary with product type, recommended ranges are provided in 5.4. If no range is provided, ambient temperature is recommended. Maintain the selected temperature range using hot trays, water baths, steam tables, crushed ice, or ice water. Ranges larger than those recommended may result in differences due to temperature rather than the test variables. The sample's temperature should not be so hot that the assessor cannot comfortably hold it in one's mouth. The temperature, however, should be selected to ensure the sample is microbiologically safe. It is important to monitor the temperature of the samples throughout serving and evaluation. Preheating serving containers will minimize heat loss. A new sample may need to be provided during a test period to maintain a consistent temperature. Also, consider the effect of temperature loss or gain on a product's moisture content, viscosity, and other characteristics.

5.1.6 *Palate Cleansers*—Palate cleansers should be used before evaluation and between samples. Different types of products require different palate cleansers. A palate cleanser needs to be as bland as possible and easily cleared from the mouth. Room temperature water, bottled, filtered, distilled, etc., is a commonly used cleanser. Saltine or oyster crackers, unsalted tops, are effective for many products. Other cleansers, such as warm water, a 50/50 blend of warm water and sodium-free carbonated water, or extended rest periods between samples also may be effective. Determine which cleanser or combination of cleansers is best for a product during pretesting.

5.1.7 *Carriers*—Consider the test objective and nature of the product when determining if a carrier is needed. Carriers most often are used to present products in a form that is more typical of how the product is used or consumed by consumers. In addition, some products may require being evaluated with a carrier, for example, fruit flavors may need to be evaluated in a sucrose/citric acid solution or a soy sauce may need to be evaluated on rice because of its high flavor intensity when evaluated plain. Product and carrier can be combined prior to serving or served separately for the assessor to combine.

5.1.7.1 Several things need to be considered when selecting a carrier. The carrier should be appropriate for the product being tested. This often includes selecting a carrier that represents one of the more common usages of the product, for example, chicken for a BBQ sauce or turkey sandwich for mayonnaise. The carrier should have a neutral influence on the evaluation of the product. The carrier should not mask product attributes critical to the evaluation, distract attention from the product, or exceed/fall short of common consumer expectations for the carrier. For example, iceberg lettuce for a pourable salad dressing evaluation or saltine crackers for a margarine evaluation are neutral, yet similar to common usage, and enable an assessor to focus on evaluating the product sample rather than the carrier. The carrier should be consistent in aroma, flavor, texture, and appearance across all presentations of the samples; any inconsistencies can bias responses to the products themselves. In addition, the same amount of carrier should be used for each sample.

5.1.7.2 It may be difficult to determine whether a carrier is the proper choice for the product evaluation. Testing both with and without the carrier can be used to determine the impact on the differentiation of samples.

5.1.7.3 Another application of carriers to be considered is as an end-use. For example, it may be appropriate to test cereal with milk added or pancakes with syrup added. Consumer research about how the product is used may help determine if a carrier is appropriate.

5.1.8 *Time Intervals*—Pretesting will determine how much time is needed to allow an assessor's senses to recover between samples. If specific time intervals are required, steps need to be taken to ensure that all assessors maintain the intervals. Timers,

stopwatches, and time-delayed computer screens may be help-ful.

5.1.9 *Expectoration*—Consider the test objective in determining if assessors need to expectorate. Differences in expectoration techniques may affect assessor results. It may be desirable to standardize the technique. Minimally, individual assessors should treat all samples consistently. Provide assessors with an appropriate and sanitary means of expectoration. Sixteen-ounce opaque cups with lids work well, since assessors can place their mouths completely and discretely in the cup opening. Assessors should be responsible for discarding their own containers.

5.1.10 *Product Holding Time*—Some products may change following preparation or when exposed to the environment so it may be important to control the time between sample preparation and serving. For example, bread and meat may dry out, candy may get tacky, soups and sauces may thicken, ice cream may melt, French fries may get soft/limp, etc. Pretest to determine the appropriate holding time.

5.1.11 *Test Environment*—The testing environment should be quiet and odor-free. Assessors and those serving the samples should keep conversation to a minimum and speak quietly when necessary. Cologne, perfume, aftershave, etc., that may interfere with the evaluation of products, should not be worn by either the serving staff or the assessors (refer to STP 913).

5.1.12 *Number of Samples*—Consider assessor fatigue, both physiological and psychological, when determining the number of samples to be evaluated in a test session. The flavor of the samples, number of questions, and length of test session should also be considered.

5.1.13 *Marking/Coding of Products and Samples*—Grease pencils or coded stickers are recommended for marking sample containers. Felt markers or pens should be used with caution due to their odor. Prepare in advance to allow the odor to dissipate.

5.2 Heated Foods Guidelines:

5.2.1 Consistency in the preparation of all samples is the primary goal. Use the appliance(s) that is commonly associated with product preparation. Consider preparing samples side-byside in the same cooking unit or balance among cooking units. All cooking units need to be routinely calibrated. Most foods should be cooked to a specific internal temperature. Some foods require heating for a specific length of time, for example, a pie needs to be heated long enough to bake the crust. The location in the sample where temperature will be taken needs to be determined and remain consistent throughout testing. The internal temperature is sometimes lower than surface temperature. Once cooked, foods should be held at a minimum of 60°C (140°F) for microbiological safety. Determine the effect of holding, or tempering, samples. This effect includes changes in the initial internal temperature, as well as other physical changes. Reheating is not recommended.

5.2.1.1 Many products do not cook uniformly, for example, a crust may form on the top and side surfaces, the outside edges may cook to a higher temperature or become browner than the interior, etc. Prior to the test, determine if only the interior portions will be used or if the entire product will be used. If the entire product is served, care must be taken to ensure that each

assessor evaluates similar portions.

5.2.1.2 Several items can be used to maintain the recommended temperature of samples prior to serving and during testing, for example, heated sand, electrical pads/plates, heat lamps, water baths, preheated glass/Pyrex[®] serving containers with lids, steam tables, etc. Equipment should be similar in terms of wattage, energy source, shelf height, etc.

5.2.2 *Microwaved*—It is recommended that the same brand, model, and wattage ovens be used for product preparation (see IEC 705). Since ovens vary in their output, even those with the same brand and model, it is recommended that samples be balanced by ovens. All ovens should be clean before each sample is placed in them. Be consistent placing products in cold or preheated ovens. Ice bags or cold towels may be appropriate for cooling ovens.

5.2.2.1 If several microwave ovens will be operating at once, they must be tested/calibrated while all are operating at the same time to be sure they are heating products similarly. Check circuit voltage to ensure uniform power supply conditions, one oven per circuit breaker is recommended. Line voltage may vary throughout the day. Extension cords are not recommended for use with microwave ovens.

5.2.2.2 Place all samples in approximately the same position in the oven. Minimize the effect of hot and cold spots by rotating samples, stirring, and using turn tables. If samples are rotated, schedule the rotation, and rotate all samples the same amount. Develop a procedure for maintaining oven temperature throughout the test. Leaving the oven door open a predetermined amount of time between heating samples allows the air in the oven to cool down. It is recommended that samples be similar in weight, size, shape, etc., to ensure even cooking, however, identical cooking time may not achieve the same end temperature. Pretesting the sample cook time is necessary.

5.2.3 *Baked/Broiled*—Standardize and monitor oven temperature throughout the test. Determine the oven temperature cycle and be consistent placing products in the ovens during that cycle. Be consistent in using electric or gas ovens for a test.

5.2.3.1 Use the proper size pans, trays, cookie sheets, etc., to ensure even cooking of the samples. It is better to have fewer samples in each oven and have them cook evenly on all sides. Allow a minimum of 2 in. between pans and oven walls for proper air circulation. Be sure that the rack is in the same position in all ovens and samples always are placed in the same position.

5.2.3.2 Toasters, toaster ovens, convection ovens, and conventional ovens heat products differently. Toasters and toaster ovens generally are not recommended as substitutes for conventional ovens. Food service equipment may be appropriate for selected applications.

5.2.4 Stove Top—Be consistent in using electric or gas burners for a test. Use uniform burner size, flame size, and heating container. Control for hot spots or variations across cooking surfaces. Heating containers vary in the way they conduct and hold heat. Consider using stainless steel double boilers for maximizing temperature maintenance over time.

5.2.5 Deep Fried—Condition fresh oil by heating oil for 1 h

at 185 to 190°C (365 to 375°F). Fry at least two preliminary batches prior to frying samples for evaluation. The recommended free fatty acid level is 1.5 %, but no more than 2 % of the U.S.D.A. maximum for processing plants. Constantly monitor the temperature of the cooking oil during a test. A hanging thermometer is ideal for this purpose. Stir the oil to ensure even heating and to help eliminate hot/cold spots before the food is put into the fryer.

5.2.5.1 When putting food into the fryer, be cautious of ice crystals causing the oil to splatter. As a general guideline, do not fill the container more than $\frac{2}{3}$ full, and maintain a constant level throughout the test. Fry under a hood, if possible, and be aware of fire hazards. Pretest to determine if food will be flipped or stirred during cooking. This ensures even frying.

5.2.5.2 Maintain a constant holding time, determined by pretesting. When holding fried foods that are coated with breading, control for moisture buildup in the food warmer.

5.2.6 *Pan Fried/Grilled*—Monitor the surface temperature in several locations of each frying pan or food service grill using a surface temperature thermometer. Calibrate the temperature dial setting of the appliance for equivalent surface temperature for all sections of the grill or each frying pan. Do not cook samples on hot or cold spots. It is recommended that the same brand and model of frying pan be used for product preparation. Be consistent placing products on cold or preheated surfaces. For products with low moisture or low fat, it may be necessary to preseason the grilling surface with oil to ensure the product does not stick to the cooking surface.

5.2.6.1 If several frying pans will be operating at once, they must be tested/calibrated while all are operating at the same time to be sure they are heating products similarly. Check circuit voltage to ensure uniform power supply conditions. Frying pans need separate circuit breakers.

5.2.6.2 When preparing samples side-by-side on the same cooking unit, take care to keep product variables separated. Construct dividers or use enough distance between product variables so that grease runoff or the product itself does not contact other products. For food service grills, establish protocol to clean and possibly reseason the grill surface between cooking samples.

5.3 Refrigerated and Frozen Foods Guidelines:

5.3.1 Foods should be held below 4°C (40°F) for microbiological safety. Control for temperature variations in the areas where samples are stored for holding, tempering, or serving. Load the storage space properly to ensure adequate air circulation and protect all samples so they do not absorb odors from equipment or other products in the storage area. Monitor the cycling patterns and resulting temperature fluctuations in these areas and the effect of these fluctuations on the samples. Set the temperature ranges for storage, tempering, and serving based on the test objective.

5.3.2 Use pretesting and temperature monitoring to determine the proper handling protocol for samples. For example, you may need to control the number of times a refrigerator or freezer is opened during a test. There are a number of alternatives for holding and serving chilled/frozen samples: insulated containers, coolers, water or ice baths, soft packs, shielded dry ice, etc. 5.3.3 Samples should be shielded from direct contact with dry ice by using air space, paper, or styrofoam as insulators. Use dry ice only in well-ventilated areas and practice adequate safety precautions to avoid burns. Discard any food that comes in direct contact with dry ice.

5.4 Specific Considerations:

5.4.1 Beverages:

5.4.1.1 *Serving Temperature*, 5 to 9° C (42 to 48° F) usually is appropriate for refrigerated beverages and 58 to 60° C (136 to 140° F) for hot beverages.

5.4.1.2 Sample Preparation:

(a) Powdered—For products served at room temperature or products that change with time, prepare and serve them immediately. For products served chilled, prepare them ahead of time and refrigerate to allow temperature to equilibrate and powder to hydrate. Reconstitute with a liquid you can control, for example, bottled water or milk. If using milk for reconstitution, select one fat level and consider combining the contents of cartons into a common lot to minimize variability.

(*b*) *Blended*—Prepare samples and serve them immediately. The pouring sequence should be consistent among samples for each assessor.

(c) Carbonated—Pour samples and serve them immediately. If using large containers, pretest to determine the maximum holding time to ensure optimal carbonation.

(d) Juice—Shake products prior to pouring samples and serve them immediately.

(e) Hot—Keep samples covered between servings. Erlenmeyer and Thermos[®] flasks are recommended for serving ease and minimal temperature loss. Flasks should be of the same volume and weight. Some sweeteners, like aspartame, are affected by heat so hold time is critical.

5.4.2 Breads/Bakery Goods/Cookies/Crackers:

5.4.2.1 Sample Preparation—Avoid variation within samples, such as edges of a cake, unless it is warranted by the test objective. If appearance is to be evaluated, the assessor should receive a whole slice of bread or a whole roll or muffin. Serve samples from the center portion of the loaves of bread to keep sample sizes uniform and avoid crusts, unless crusts should be included as determined by the test objective. Consider whether it is appropriate to serve rolls baked in the center or on the edges of the baking band, or both, since those baked on the edges may be darker in color. Ambient samples may need to be placed in closed containers during the test to prevent changes in moisture content.

5.4.2.2 For cookies and crackers, consider serving edge portions, as well as interior portions as determined by the test objective. Prior to the test, decide if over-browned samples will be used. Serve products right side up and do not serve broken pieces. These products may need to be kept in closed plastic bags during the test to prevent changes in moisture content.

5.4.3 Cereals:

5.4.3.1 *Serving Temperature*—Ambient for RTE (ready to eat) cereal, 43 to 49°C (110 to 120°F) for hot cereal.

5.4.3.2 *Holding Time*—Determine the maximum sample holding time for hot cereal by pretesting.

5.4.3.3 Sample Preparation—Discrimination and descriptive tests may require counting of additives, for example, granola, fruits, marshmallows, nuts, etc. For affective tests, cereal is evaluated commonly as it comes from the box in any proportion ensuring that reasonable consistency is maintained. Samples should be served from the top two thirds of the box to avoid fines.

5.4.3.4 Milk and sugar often are used in affective tests. Assessors typically use the type and amount they prefer. If used in discrimination and descriptive tests, however, all assessors should use the same amount of the same type. Consider the synergistic effect of artificial sweeteners when determining if they should be used.

5.4.4 Confections:

5.4.4.1 *Chocolate Bars*—If bars are stamped with a brand name, consider shaving off the name or chopping the bars into smaller pieces so the name is not identifiable.

5.4.4.2 *Mints/Hard Candies*—Pretest to determine how long the assessor should hold the sample in one's mouth before evaluating. Forty-five to sixty seconds usually is appropriate with a minimum of a 2 to 3-min interval between samples.

5.4.4.3 *Marshmallows/Chewy Candies/Chewing Gum*—Pretest to determine the length of chewing before evaluation(s).

5.4.5 Dairy/Dairy-Type Products:

5.4.5.1 *Block Cheese*—Serving temperature should be ambient or refrigerated depending on the test objective. Trim the outer portion of block cheeses before preparing samples. Cheese wire or nylon thread may be useful for cutting soft cheeses. Remove wrappers from prewrapped sliced cheese and trim the edges, unless assessors need to evaluate if cheese sticks to wrapper. Control the time between preparing cheese samples and serving them to minimize changes in moisture content. Keep samples covered with plastic wrap and store in the refrigerator if necessary.

5.4.5.2 *Cottage Cheese/Cream Cheese/Sour Cream/Yogurt*: (*a*) *Serving Temperature*—Ambient or refrigerated depending on the test objective.

(b) Sample Preparation—Prior to the test, determine if any liquid resulting from syneresis will be reincorporated into the samples. Scoop from the same area of the container. Pretest scooping procedures to ensure sample uniformity. Consider stirring yogurt prior to preparing samples if products have fruit at the bottom.

(c) Carriers—Consider room-temperature bagels as carriers for cream cheese and baked potatoes for sour cream.

5.4.5.3 *Fluid Milk*—Serving temperature usually is refrigerated. Protect samples from exposure to light to prevent oxidation. Foil-wrapped containers or tinted glass are effective for protecting samples.

5.4.6 *Eggs*:

5.4.6.1 Direct heat is not recommended when preparing scrambled eggs for evaluation. Consider cooking eggs in a double boiler over hot water. Cook the eggs without adding salt, margarine, etc., to avoid confounding the variables being evaluated.

5.4.6.2 End-point cooking temperatures should range from 76 to 78°C (170 to 174°F). Eggs cool and dry out quickly so they should be served immediately after cooking. If it is necessary to hold cooked eggs, however, consider putting them

in warmed jars with lids, such as baby food jars, and placing them in warmed sand held in a warming cabinet or oven.

5.4.7 Entrees:

5.4.7.1 *Serving Temperature*—60 to 71°C (140 to 160°F) usually is appropriate.

5.4.7.2 *Serving Size*—Product integrity is an important consideration when determining serving size, for example, portions should not be used that adversely affect the product's appearance, moisture content, physical properties, etc. Depending on the test objective, it may be necessary to have assessors evaluate the appearance of the entire entree in its original packaging and then an individual serving in a test container.

5.4.8 Fresh Produce:

5.4.8.1 *Serving Temperature*—Ambient or refrigerated depending on the test objective.

5.4.8.2 Sample Preparation—Use nonmetal utensils to prepare samples to reduce browning. Prepare samples within 1 h of serving to maintain fresh appearance or consider having the assessors cut their own samples. Provide knives and cutting boards and instruct the assessors on how to cut the produce. Consider serving two to three pieces of each sample as produce varies from root to root and plant to plant. Assessors should evaluate portions of the entire sample.

5.4.9 Frozen Dessert Products:

5.4.9.1 Serving Temperature— -18 to -10° C (0 to 14° F). After samples are prepared for a test, retemper them to a uniform serving temperature.

5.4.9.2 *Time Intervals*—Keep the number of samples evaluated in a session to a minimum, and allow enough time between samples for the assessor's mouth to return to normal body temperature.

5.4.9.3 *Sample Storage*—Bulk frozen desserts should temper until a consistent temperature suitable for sample preparation is reached: -18 to -10° C (0 to 14° F) overnight is recommended. Place products side by side in the freezer to minimize temperature gradients within the box.

5.4.9.4 Sample Preparation-Scoop from the same area of the carton. Pretest scooping procedures to ensure sample uniformity. Avoid scraping the sides as that product tends to lose its original identity. Consider slicing product packed in cartons for easier preparation than scooping. Consider cutting bars into smaller portions depending on the test objective and whether or not the product has a stick. Be sure that each assessor gets a sample from the same location of the container for all products. Cut coated products prior to tempering to reduce the possibility that the coating structure will collapse. Minimize temperature fluctuations during the test. Consider using portable chest freezers to store products during the test as vertical freezers tend to allow the cold air to spill out when the door is opened. This can be controlled by taping plastic over the face of the door and cutting holes in the plastic for reaching into the freezer. If more than one freezer is needed, consider rotating samples in freezers for holding and serving. Serve samples immediately after removing from tempering freezer.

5.4.10 Jams/Preserves/Jellies/Fruit Spreads:

5.4.10.1 *Serving Temperature*—Ambient or refrigerated depending on the test objective.

5.4.10.2 *Sample Preparation*—Serve a sample that is representative of the product in the consumer package, for example, consider fruit pieces, as well as jelled portion.

5.4.10.3 *Carriers*—Consider white bread or unsalted crackers as carriers. A firm-textured white bread is useful for assessing spreadability. Use a nonserrated knife or back of a spoon for spreading.

5.4.11 Margarine/Butter/Spreads:

5.4.11.1 *Serving Temperature*—Ambient or refrigerated depending on the test objective.

5.4.11.2 *Carriers*—Consider white bread or unsalted crackers as carriers. A firm-textured white bread is useful for assessing spreadability. Use a nonserrated knife or back of a spoon for spreading.

5.4.12 *Meats/Fish/Poultry*—Cook products to a minimal internal temperature to destroy bacteria. The following temperatures are the minimum needed to destroy bacteria; however, some products may need to be cooked to a higher temperature for product palatability. Pretest to determine the appropriate temperature.

5.4.12.1 Beef Patties: 71°C (160°F).

5.4.12.2 Beef Steaks/Roasts: 63°C (145°F).

5.4.12.3 Pork Chops/Roasts: 63°C (145°F).

5.4.12.4 Fish: 66°C (150°F).

5.4.12.5 Poultry Patties: 74°C (165°F).

5.4.12.6 Poultry Breast: 63°C (145°F).

5.4.12.7 Serve at 60°C (140°F).

5.4.12.8 Cold items, such as cold cuts, lunch meats, etc., should be stored and served between 2 and 5°C (35 and 40°F). Do not serve above 7°C ($45^{\circ}F$) to ensure samples are microbiologically safe. Samples should be similar in size, shape, and physical condition, that is, muscle configuration, etc. For further information about meat, fish, and poultry cooking and preparation, see Refs (1-12).

5.4.13 Oils/Fats-Refer to Practice E 1627.

5.4.14 *Pancakes/Waffles from Batter*—Sample preparation includes mixing the batter consistently each time. Consider using an automatic mixer with a whip attachment. Pretest to

determine if batter needs to sit before pouring on the grill. Use a standard scoop to ensure the same sized samples and monitor the batter temperature throughout the test. Discard atypical samples, due to incorrect flipping, folding, etc. Use a seasoned grill. Oil should not be added to the grill during the testing. Serve samples immediately after cooking.

5.4.15 Pourable Salad Dressings:

5.4.15.1 *Serving Temperature*—Ambient or refrigerated depending on the test objective.

5.4.15.2 *Sample Preparation*—The pouring sequence should be consistent among samples for each assessor.

5.4.15.3 *Carriers*—Consider chopped, well-dried iceberg lettuce or prepackaged lettuce as a carrier.

5.4.16 Soups/Sauces/Gravies:

5.4.16.1 *Serving Temperature*—60 to 71°C (140 to 160°F) usually is appropriate.

5.4.16.2 *Carriers*—Consider instant mashed potatoes as a carrier for gravies, unsalted tortilla chips for cheese sauces, and plain boiled rice for strongly flavored sauces.

5.4.17 Snack Foods:

5.4.17.1 *Snack Cakes*—Remove wrappers for serving, taking care to not alter the appearance.

5.4.17.2 *Snack Mixes*—Consider counting the additives, depending on the test objective.

5.4.17.3 *Potato Chips/Pretzels*—Serve only whole chips or pretzels or consider breaking large pieces into smaller ones if uniform appearance is needed for the test objective.

5.4.18 *Syrups*:

5.4.18.1 *Serving Temperature*—Ambient, refrigerated, or heated depending on the test objective.

5.4.18.2 *Carriers*—Consider pancakes or waffles as carriers for breakfast syrups (baked mini waffles and frozen pancakes are convenient. Consider milk or vanilla ice cream for other syrups, such as chocolate or strawberry.

6. Keywords

6.1 foods and beverages; sensory evaluation; serving protocol

REFERENCES

- (1) AMSA 1995. "Research Guidelines for Cookery, Sensory Evaluation and Instrumental Tenderness Measurements of Fresh Meat." American Meat Science Association, Chicago, IL.
- (2) ASTM Committee E-18, 1996. "Sensory Testing Methods, 2nd Edition, Manual 26," American Society for Testing Materials, West Conshoshocken, PA.
- (3) ASTM Committee E-18 Guidelines for the Selection and Training of Sensory Panel Members; STP 758.
- (4) ASTM Committee E-18 Physical Requirements for Sensory Evaluation Laboratories; STP 913.
- (5) Beery, K. E., and Ziegler, J. H., "A simplified and rapid method to uniformly size rib steak pieces for taste panel evaluation," *J. Food Science*, Vol 34, 1969, p. 480.
- (6) Campbell, A. M., Penfield, M. P., and Griswold, R. M., The experimental study of food." 2nd ed., Houghton Mifflin, Boston, MA, 1979.

- (7) Caporaso, F. "A simple technique for maintaining temperature of cooked meat samples prior to sensory evaluation." *J. Food Science*, Vol 43, 1978, p. 1041.
- (8) Carlin, A., FD., and Harrison, D. L., "Cooling and sensory methods used in experimental studies on meat." National Live Stock and Meat Board, Chicago, IL, 1978.
- (9) FSIS Directive 7370.2 1995. United States Department of Agriculture, Food Safety and Inspection Service, Washington, DC.
- (10) Poste, L. M., Mackie, D. A., Butler, G., and Larmond, E. "Laboratory methods for sensory evaluation of food." Publication 1864/E, Canada Department of Agriculture, Ottawa, 1991.
- (11) USDA Meat and Poultry Hotline 1-800-535-4555.
- (12) Warren, M. W., and Ball, H. R., "Effect of concentration of egg yolk and white on fresh scrambled egg flavor." *Poultry Science*, Vol 70, 1991, pp. 2186–2190.

The American Society for Testing and Materials takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, 100 Barr Harbor Drive, West Conshohocken, PA 19428.

This standard is copyrighted by ASTM, 100 Barr Harbor Drive, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (http://www.astm.org).